Please check the examination details below before entering your candidate information

Candidate surname	Other names

Centre Number | Candidate Number

October 2021

Pearson Edexcel International Advanced Level

Time 1 hour 30 minutes

Paper reference **WME01/01**

Mathematics

International Advanced Subsidiary/Advanced Level
Mechanics M1

You must have:
Mathematical Formulae and Statistical Tables (Yellow), calculator

Total Marks

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided – *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Whenever a numerical value of g is required, take $g = 9.8\,\text{m s}^{-2}$, and give your answer to either 2 significant figures or 3 significant figures.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 8 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets – *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over ▶

P66649A

E:1/1/1/

P 6 6 6 4 9 A 0 1 3 2

Pearson

Leave blank

1.



Figure 1

A non-uniform rod AB has length 9 m and mass M kg.

The rod rests in equilibrium in a horizontal position on two supports, one at C where $AC = 2.5$ m and the other at D where $DB = 2$ m, as shown in Figure 1.

The magnitude of the force acting on the rod at D is twice the magnitude of the force acting on the rod at C.

The centre of mass of the rod is d metres from A.

Find the value of d.

(6)

Question 1 continued

Leave blank

Leave blank

Question 1 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Leave blank

Question 1 continued

Q1

(Total 6 marks)

Leave blank

2. A particle P of mass $2m$ is moving on a rough horizontal plane when it collides directly with a particle Q of mass $4m$ which is at rest on the plane. The speed of P immediately before the collision is $3u$. The speed of Q immediately after the collision is $2u$.

(a) Find, in terms of u, the speed of P immediately after the collision.

(3)

(b) State clearly the direction of motion of P immediately after the collision.

(1)

Following the collision, Q comes to rest after travelling a distance $\frac{6u^2}{g}$ along the plane.

The coefficient of friction between Q and the plane is μ.

(c) Find the value of μ.

(6)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Leave blank

Question 2 continued

Leave blank

Question 2 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Leave blank

Question 2 continued

Q2

(Total 10 marks)

Leave blank

3. A car is moving at a constant speed of $25\,m\,s^{-1}$ along a straight horizontal road.

The car is modelled as a particle.

At time $t = 0$, the car is at the point A and the driver sees a road sign 48 m ahead.

Let t seconds be the time that elapses after the car passes A.

In a **first** model, the car is assumed to decelerate uniformly at $6\,m\,s^{-2}$ from A until the car reaches the road sign.

(a) Use this first model to find the speed of the car as it reaches the sign.

(2)

The road sign indicates that the speed limit immediately after the sign is $13\,m\,s^{-1}$.

In a **second** model, the car is assumed to decelerate uniformly at $6\,m\,s^{-2}$ from A until it reaches a speed of $13\,m\,s^{-1}$. The car then maintains this speed until it reaches the road sign.

(b) Use this second model to find the value of t at which the car reaches the sign.

(4)

In a **third** model, the car is assumed to move with constant speed $25\,m\,s^{-1}$ from A until time $t = 0.2$, the car then decelerates uniformly at $6\,m\,s^{-2}$ until it reaches a speed of $13\,m\,s^{-1}$. The car then maintains this speed until it reaches the road sign.

(c) Use this third model to find the value of t at which the car reaches the sign.

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Leave blank

Question 3 continued

Leave blank

Question 3 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Leave blank

Question 3 continued

(Total 10 marks)

Q3

Leave blank

4. The position vector, **r** metres, of a particle P at time t seconds, relative to a fixed origin O, is given by

$$\mathbf{r} = (t - 3)\mathbf{i} + (1 - 2t)\mathbf{j}$$

(a) Find, to the nearest degree, the size of the angle between **r** and the vector **j**, when $t = 2$

(3)

(b) Find the values of t for which the distance of P from O is 2.5 m.

(5)

Leave blank

Question 4 continued

Leave blank

Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Leave blank

Question 4 continued

Q4

(Total 8 marks)

Leave blank

5.



Figure 2

A small bead of mass 0.2 kg is attached to the end P of a light rod PQ. The bead is threaded onto a fixed vertical rough wire.

The bead is held in equilibrium with the rod PQ inclined to the wire at an angle α, where $\tan\alpha = \frac{4}{3}$, as shown in Figure 2.

The thrust in the rod is T newtons.

The bead is modelled as a particle.

(a) Find the magnitude and direction of the friction force acting on the bead when $T = 2.5$

(3)

The coefficient of friction between the bead and the wire is μ.

Given that the greatest possible value of T is 6.125

(b) find the value of μ.

(7)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 5 continued

Leave blank

Leave blank

Question 5 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Leave blank

Question 5 continued

Q5

(Total 10 marks)

Leave blank

6.



Figure 3

A small ball is thrown vertically upwards at time $t = 0$ from a point A which is above horizontal ground. The ball hits the ground 7 s later.

The ball is modelled as a particle moving freely under gravity.

The velocity-time graph shown in Figure 3 represents the motion of the ball for $0 \leqslant t \leqslant 7$

(a) Find the speed with which the ball is thrown.

(2)

(b) Find the height of A above the ground.

(3)

Leave blank

Question 6 continued

Q6

(Total 5 marks)

Leave blank

7.



Figure 4

One end of a light inextensible string is attached to a particle A of mass $2m$. The other end of the string is attached to a particle B of mass $3m$. The string passes over a small, smooth, light pulley P which is fixed at the top of a rough inclined plane. The plane is inclined to the horizontal at an angle α, where $\tan\alpha = \frac{3}{4}$

Particle A is held at rest on the plane with the string taut and B hanging freely below P, as shown in Figure 4. The section of the string AP is parallel to a line of greatest slope of the plane.

The coefficient of friction between A and the plane is $\frac{1}{2}$

Particle A is released and begins to move up the plane.

For the motion before A reaches the pulley,

(a) (i) write down an equation of motion for A,

(ii) write down an equation of motion for B,

(4)

(b) find, in terms of g, the acceleration of A,

(5)

(c) find the magnitude of the force exerted on the pulley by the string.

(4)

(d) State how you have used the information that P is a smooth pulley.

(1)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Leave blank

Question 7 continued

Leave blank

Question 7 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Leave blank

Question 7 continued

(Total 14 marks)

Q7

Leave blank

8. [*In this question* **i** *and* **j** *are horizontal unit vectors directed due east and due north respectively and position vectors are given relative to a fixed origin.*]

At 7 am a ship leaves a port and moves with constant velocity. The position vector of the port is $(-2\mathbf{i} + 9\mathbf{j})$ km.

At 7.36 am the ship is at the point with position vector $(4\mathbf{i} + 6\mathbf{j})$ km.

(a) Show that the velocity of the ship is $(10\mathbf{i} - 5\mathbf{j})\,\text{km h}^{-1}$

(2)

(b) Find the position vector of the ship t hours after leaving port.

(2)

At 8.48 am a passenger on the ship notices that a lighthouse is due east of the ship.

At 9 am the same passenger notices that the lighthouse is now north east of the ship.

(c) Find the position vector of the lighthouse.

(4)

(d) Find the position vector of the ship when it is due south of the lighthouse.

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Leave blank

Question 8 continued

Leave blank

Question 8 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 8 continued

Leave blank

Leave blank

Question 8 continued

Q8

(Total 12 marks)

TOTAL FOR PAPER: 75 MARKS

END

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA